

The Best of


# EXOTIC




**DVM**  
VOLUME 12  
ISSUE 4

*Selected Practical Peer-reviewed Papers for Clinicians*

## Exotic Animal Medicine & Surgery



*Contents*

- 5** **Extraction of Cheek Teeth and Surgical Treatment of Periodontal Abscessation in Pet Rabbits with Acquired Dental Disease**  
Vittorio Capello, DVM, Dip ECZM (Small Mammal), Dipl ABVP (ECM)
- 13** **Fluid Resuscitation and Nutritional Support in Rabbits with Gastric Stasis or Gastrointestinal Obstruction**  
Marla Lichtenberger, DVM, Dipl ACVECC
- 18** **Ferret Adrenal Removal Using Temporary Occlusion of the Caudal Vena Cava**  
R. Avery Bennett, DVM, MS, Dipl ACVS
- 21** **Clinical Aspects of Inflammatory Bowel Disease in Ferrets**  
Mark Burgess, DVM and Michael M. Garner, DVM, Dipl ACVP
- 27** **What Veterinarians Need to Know About Guinea Pigs**  
Cathy A. Johnson-Delaney, DVM, Dipl ABVP (Avian), Dipl ABVP (ECM)
- 34** **Flank Approach to Elective Ovariectomy in Guinea Pigs**  
Vittorio Capello, DVM, Dip ECZM (Small Mammal), Dipl ABVP (ECM)
- 37** **Surgical Techniques in Pet Hamsters**  
Vittorio Capello, DVM, Dip ECZM (Small Mammal), Dipl ABVP (ECM)
- 43** **Application of Diagnostic Endoscopy in Birds**  
Michael Lierz, Dr med vet, MRCVS, DZooMed, Dip ECZM (Wildlife Population Health), Dip ECPVS
- 49** **Minimally Invasive Endosurgery of the Psittacine Cranial Coelom**  
Stephen J. Divers, BVetMed, DZooMed, Dipl ACZM, Dip ECZM (Herpetology), FRCVS; Michael McBride, DVM; Chris Hanley, DVM, Dipl ACZM and Heather Barron, DVM, Dipl ABVP (Avian)
- 54** **Surgical Approach to the Thoracic Cavity of Birds**  
R. Avery Bennett, DVM, MS, Dipl ACVS
- 57** **Ventral Midline Approach to Avian Salpingohysterectomy**  
Marc H. Kramer, DVM and Don J. Harris, DVM
- 62** **Surgical Resolution of Reproductive Disorders in Female Green Iguanas**  
Scott J. Stahl, DVM, Dipl ABVP (Avian)
- 67** **Endoscopic Evaluation of the Reptilian Respiratory System**  
Stephen J. Divers, BVetMed, DZooMed, Dipl ACZM, Dip ECZM (Herpetology), FRCVS; Michael McBride, DVM; Chris Hanley, DVM, Dipl ACZM and Heather Barron, DVM, Dipl ABVP (Avian)
- 71** **Flap Closure Method Using Epoxy Putty in Plastron Osteotomy in Chelonians**  
Kenichi Tamukai, DVM
- 78** **5-Minute Guide to Amphibian Disease**  
Mads F. Bertelsen, DVM, DVSc, Dipl ACZM, Dip ECZM (Herpetology) and Graham Crawshaw, BVetMed, MRCVS, Dipl ACZM
- 82** **Practical Koi and Goldfish Medicine**  
Dan H. Johnson, DVM, Dipl ABVP (ECM)
- 89** **Introduction to Invertebrate Medicine**  
Gregory A. Lewbart, MS, VMD, Dipl ACZM



# Surgical Techniques in Pet Hamsters

Vittorio Capello, DVM, Dip ECZM (Small Mammal),  
Dipl ABVP (ECM)



Vittorio Capello, DVM, Dip ECZM  
(Small Mammal), Dipl ABVP (ECM)  
capellov@tin.it

**Table 1. Anesthetic and Analgesic Agents in Hamsters**


<b>Induction</b>	<ul style="list-style-type: none"> <li>• <b>Isoflurane</b> - 3-5% in oxygen (mask directly for patients at increased anesthetic risk or for Russian hamsters)</li> <li>• <b>Ketamine + xylazine</b> - ketamine (3-5 mg/100 g IM) + xylazine (0.3-0.5 mg/100 g IM)</li> <li>• <b>Ketamine + medetomidine</b> - ketamine (3-5 mg/100 g IM) + medetomidine (7-10 µg/100 g IM)</li> </ul>
<b>Maintenance</b>	<ul style="list-style-type: none"> <li>• <b>Isoflurane</b> - 1-3% in oxygen; delivered by facemask</li> </ul>
<b>Analgesia</b>	<ul style="list-style-type: none"> <li>• <b>Butorphanol</b> - 0.05-0.1 mg/100 g SC q4h</li> </ul>

Note: Particular attention must be paid to proper dilution of all anesthetic and analgesic agents.



## Anesthesia



**Fig 1.** Facemasks used for hamster anesthesia should be transparent so the mucous membrane color of the nose and tongue can be easily checked. For the very small Russian hamsters (body weight ~28 g), a properly cut and contoured syringe may be used as a facemask.



**Fig 2.** Cardiac activity can be monitored in anesthetized hamsters with a vascular Doppler system. The probe is positioned on the chest and held in place by the anesthetist or by the adhesive drape. Rectal temperature is also monitored by a very small probe. In the author's experience, pulse oximetry is difficult to apply to the hamster. Transparent or semi-transparent drapes are mandatory for hamster surgery to allow for monitoring respiration.



**Fig 3. a)** Fluids may be administered through the intraosseous (IO) route by inserting a 25-ga (or smaller) needle into the medullary cavity of the tibia. **b)** Special attention must be paid during insertion, and the position of the needle should be checked with radiographs in both the lateral and sagittal projections.


Originally published in *Exotic DVM* Volume 5.3, pp 19-24

## Laparotomy


Extraordinary care must be taken with anesthesia and basic surgical principles when performing a celiotomy in a hamster (Fig 4). Heating devices are mandatory. If possible, intraosseous fluids should be administered to the patient. Fur should be shaved to create a wide surgical field, and transparent adhesive drapes are recommended (Figs 5-8).

The abdominal wall is closed with 5-0 to 6-0 absorbable monofilament suture. A simple continuous suture pattern reduces surgical time, but a simple interrupted pattern can be used instead. The subcutis does not need to be sutured in a separate layer. The skin is sutured with 5-0 to 6-0 absorbable or non-absorbable material (e.g., nylon) in a simple interrupted pattern.


Most hamsters, except those with very calm behavior, will attempt to remove skin sutures, and an Elizabethan collar is not an option. Therefore, 5-0 absorbable monofilament suture is used in a subcuticular pattern, and tissue glue is applied to the skin incision.



**Fig 4.** Gross anatomy of abdominal organs as they appear during celiotomy. The large cecum lies immediately beneath the linea alba; however, it is easy to see because of its greenish color. (Modified from Popesko.<sup>14</sup>)



**Fig 5.** Retraction of the abdominal wall with forceps along the linea alba will prevent unintentional incision of the cecum. During surgery, abdominal tissues and organs should be kept moist by instilling warm saline into the abdomen. Retraction of the abdominal wall after the incisions may be performed using an eyelid retractor. The Lone Star Retractor may also be applied with golden hamsters if applied with care.



**Fig 6.** A large hepatic cyst is detected and isolated during explorative laparotomy in a 2-year-old female golden hamster. The cyst was attached to the liver by a thin strip of tissue. The stalk was ligated, and the entire cyst was removed.



**Fig 7.** Hemangiosarcoma was found in the spleen of this Russian hamster.



**Fig 8.** The testicle in this Russian hamster is abscessed.

## Ovariohysterectomy


Therapeutic ovariohysterectomy is indicated in case of ovarian cysts, endometritis or uterine neoplasia (Figs 9,10). After the abdominal wall is incised, the cecum must be gently retracted to expose the uterine horns. In cases of septic endometritis or uterine neoplasia, the horns can appear immediately under the linea alba. (See Exotic DVM Volume 5.4, pp 22-24 for a step-by-step ovariohysterectomy procedure.)

## Orchiectomy

The primary therapeutic indications for orchiectomy in hamsters are abscessation or neoplasia involving the testicle or epididymis (Fig 11). The author prefers to perform bilateral orchiectomy, even if only one testicle is affected. (See Exotic DVM Volume 5.4, pp 25-26 for a step-by-step orchiectomy procedure.)

## Excision of Skin and Other Soft Tissue Tumors


Skin and soft tissue tumors are common in hamsters. Even large tumors can be excised due to the hamster's loose subcutaneous tissue. The prognosis for surgery is related to the hamster's age and condition as well as characteristics of the tumor, such as position, extension, vascular supply and adhesion to surrounding tissues (Figs 12,13).




**Fig 9.** This sonogram from a 17-month-old golden hamster reveals an ovarian cyst, which is caudal to, and nearly as large as, the kidney.



**Fig 10.** A 2-year-old golden hamster was presented for anorexia and hunched posture. **a)** Bloody vaginal discharge was evident on physical examination, and palpation of the abdomen revealed a rounded mass, which was easily detected on radiographs (arrow).



**b)** During laparotomy, a large uterine mass was encountered (b). The uterine horns were larger than normal, and a large cyst was present in the left ovary. Ovariohysterectomy was performed. The histopathologic diagnosis was a uterine fibroma.



**Fig 11.** The epididymis in this 13-month-old golden hamster is abscessed.



**Fig 12.** A Russian hamster is anesthetized for excision of a large skin mass.



**Fig 13.** This 8-month-old golden hamster had a sarcoma of the scrotal skin.

## Surgery of Cheek Pouches

Diseases of the cheek pouches include impaction, prolapse, abscessation and neoplasia.

Impaction is often the result of feeding inadequate or inappropriate food (such as chocolate or extremely large or small seeds) or using cotton, paper or other improper bedding materials. Sometimes homolateral forelimb agenesis or amputation can

lead to impaction, because the hamster cannot adequately empty the cheek pouch. Prolapse and abscessation are common sequelae to impaction.

Cheek pouch prolapse, due to overfeeding, is relatively common in Russian hamsters. Retained food becomes adherent to the mucosal lining of the cheek pouch, and when the hamster attempts to empty it, the pouch prolapses along with the food. If the prolapse is recent and there are no

lesions on the mucosal surface, repositioning can be attempted (Figs 14-17).

Cheek pouch amputation is considered in cases of severe mucosal lesions, recurrent pouch prolapse or neoplastic disease. After surgery, it is impossible to prevent the hamster from breaking down sutures by trying to fill the cheek pouch, so the author tube feeds formula (Critical Care for Herbivores) and removes all other food and bedding for 24-36 hours (Figs 18-20).



**Fig 14.** The right cheek pouch is prolapsed in this Russian hamster.



**Fig 15** The mucosal surface is lubricated, and the pouch is gently replaced using cotton swabs.



**Fig 16.** A suture transfixes the skin and the cheek pouch mucosa to prevent relapse.



**Fig 17.** This is the same hamster 2 days later. The transfixing suture can remain in place up to 10-12 days to ensure prevention of relapse.



**Fig 18.** In another case, the cheek pouch is completely everted. A hemostat is placed at the base of the pouch, and the tissue is transected with a blade or scissors.



**Fig 19.** A simple interrupted suture is placed using absorbable 5-0 material.



**Figs 20.** **a)** This hamster had a sarcoma of the cheek pouch. **b)** The same hamster is shown 10 days after cheek pouch amputation.

## Abscesses

Similar to rabbits, treatment of abscesses requires total excision including the abscess capsule. Antibiotic therapy depends on culture and sensitivity results from the capsule wall (Fig 21).

## Enucleation

Enucleation is indicated for cases of severe monolateral panophthalmitis caused by traumatic injuries, bites or bacterial infection (Figs 22-23). If infection is suspected, systemic antibiotics and local therapy should be administered for 5-7 days prior to performing surgery. In the author's experience with some cases of pasteurellosis, bilateral panophthalmitis can occur. If clinical conditions are stable, surgery might be delayed for several days in order to evaluate the integrity of the contralateral eye.

Depending on the condition of the eyelids, enucleation can be performed

using a transpalpebral or transconjunctival technique. Using the latter technique, eyelids can be surgically debrided and sutured using 4-0 to 6-0 absorbable suture in a continuous or interrupted pattern. Conjunctival mucosa, sclera and external muscles of the globe are bluntly dissected. Hemorrhage is controlled by ligation, norepinephrine-moistened cotton swabs or a radiosurgical unit.

## Rectal Prolapse/Intussusception

This is a surgical condition, because the prolapse (Fig 24a,b) follows bowel intussusception (Fig 24c). It is mandatory that a laparotomy be performed including a reduction or resection of the prolapsed intestine and intestinal plication to prevent recurrent prolapse. Unless the problem is diagnosed very early, prognosis is poor.

## Intramedullary Pinning of Tibial Fractures

Tibial fractures occur quite frequently in hamsters often due to entrapment of the foot in the cage or wheel bars. In the author's experience, splinting does not promote bone healing. Moreover, the typical distal metaphyseal fracture is often exposed. Intramedullary (IM) pinning is the simplest and most practical method of fixation (except for comminuted fractures), even if a slight rotation of the foot cannot always be prevented (Figs 25, 26).

## Hindlimb Amputation

Hindlimb amputation is indicated in case of severe fractures, foot necrosis due to constriction injuries (e.g., cotton threads), severe abscessation due to bite wounds and neoplasia. Amputation can be performed at different levels: hip, stifle joint or midshaft femur or tibia (Fig 27).



**Fig 21.** An abscess of the ear canal in this Russian hamster is being excised.




**Fig 22.** This 12-month-old male golden hamster has severe panophthalmitis involving the left globe.




**Fig 23.** The eyelids have been sutured after enucleation using the transconjunctival technique.




**Fig 24 a-c.** Prolapse of the rectum (a) or colon (b) and bowel intussusception (c) represent severe pathology in the golden hamster. These conditions are predisposed by intestinal parasitism, proliferative ileitis ("wet tail") and diarrhea. The author has not seen rectal prolapse in Russian hamsters.



**Figs 25 a,b.** The radiographs show a midshaft tibial fracture in a golden hamster before (a) and after (b) IM pinning.



**Figs 26 a-e.** The needle from a 22-gauge intravenous catheter becomes the IM pin used to repair this tibial fracture in a hamster (a). The fracture site is exposed, and the IM pin is checked for size against the medullary cavity of the distal bone fragment (b). Another pin is inserted from the fracture site retrograde through the proximal bone fragment to trespass in the proximal tibial surface and is then removed (c). The IM pin is inserted normograde through the hole previously created in the proximal tibia (d). The fracture is reduced and stabilized by gently driving the IM pin into the distal bone fragment (e).



**Fig 27 a-c.** A 21-month-old female golden hamster was presented for severe lameness and a large mass involving the left hind limb (a). Amputation of the entire limb was performed at the level of the hip as indicated in this radiograph (b). Histopathologic examination revealed a soft tissue fibrosarcoma. At the time of follow-up, the hamster walked without discomfort, leaning on the base of the tail (c).

## Acknowledgment

The author thanks Germana Scerbanenco, DVM for contributions to this article.

## References and Further Reading

1. Anderson NL: Basic husbandry and medicine of pocket pets. *In* Birchard SJ, Sherding RG (eds): *Saunders Manual of Small Animal Practice*. Philadelphia, WB Saunders, 1994, pp 1363-1389.
2. Anderson NL: Intraosseous fluid therapy in small exotic animals. *In* Bonagura JD (ed): *Kirk's Current Veterinary Therapy XII*. Philadelphia, WB Saunders, 1995, pp 1331-1335.
3. Bauck LA, Hagan RJ: Cystotomy for treatment of urolithiasis in a hamster. *J Am Vet Med Assoc* 184:99-100, 1984.
4. Brown SA, Rosenthal KL: *Self-Assessment Colour Review of Small Mammals*. Manson Publishing, 1997.
5. Capello V: Pet hamster medicine and surgery, part III: Infectious, parasitic and metabolic diseases. *Exotic DVM* 3(6):27-32, 2002.
6. Carpenter JW: *Exotic Animal Formulary 3rd ed*, Saunders, 2004.
7. Donnelly TM: Disease problems of small rodents. *In* Hillyer EV, Quesenberry KE (eds): *Ferrets, Rabbits and Rodents: Clinical Medicine and Surgery*. Philadelphia, WB Saunders Co, 1997, pp 307-327.
8. Harkness JE, Wagner JE: *The Biology and Medicine of Rabbits and Rodents 4th ed*. Philadelphia, Williams & Wilkins, 1995.
9. Johnson-Delaney C: Ovariohysterectomy in a rat. *Exotic DVM* 4(4):17-21, 2002.
10. Leck S: What every veterinarian needs to know about hamsters. *Exotic DVM* 2(5):38-41, 2000.
11. Mason DE: Anesthesia, analgesia and sedation for small mammals. *In* Hillyer EV, Quesenberry KE (eds): *Ferrets, Rabbits and Rodents: Clinical Medicine and Surgery*. Philadelphia, WB Saunders Co, 1997, pp 378-391.
12. Mullen H: Soft tissue surgery. *In* Hillyer EV, Quesenberry KE (eds): *Ferrets, Rabbits and Rodents: Clinical Medicine and Surgery*. Philadelphia, WB Saunders Co, 1997, pp 329-336.
13. Percy DH, Barthold SW: Hamster. *In* *Pathology of Laboratory Rodents and Rabbits 3rd ed*. Blackwell, 2007, pp 179-205.
14. Popesko P, Rajtová V, Horák J: *A Colour Atlas of Anatomy of Small Laboratory Animals Vol 2: Rat, Mouse, Hamster*. Bratislava, Wolfe Publishing Ltd, 1992.

